



Affinity Bio Technologies

Quality, Experience, Passion

Affinity Complex Boron 15%

A complexed boric acid for improved boron uptake, translocation and metabolism.

CAUTION:

Avoid contact with eyes. Keep out of reach of children.
If swallowed seek medical advice.

Do not store in direct sunlight.

CONDITIONS OF SALE:

Affinity products are manufactured to label specifications.
Affinity takes no responsibility for product damage, loss,
injury or misuse of the product.

COMPATIBILITY

Affinity Complex Boron is compatible in most ag chem tank
mixes. However a jar test is recommended as some
chemicals can react. Do not mix amines or calcium products.

Do not store mixed product. Mix well prior to use.
Refer to SDS for correct handling and safety protocol.

Affinity Liquid Fertilisers works best when plants are
**actively growing and under reduced stress. Apply mid
morning in dry conditions for best results. Do not spray
in hot conditions. For best results consult a certified
agronomist to determine crop deficiencies.**

ANALYSIS

Boron..... 15%
Nitrogen..... 6%

Affinity Liquid Fertilisers

Phone (08) 8267 4774

www.affinitybiotech.com.au

DO NOT EXCEED RECOMMENDED APPLICATION RATES

Foliar Applications:

Multiple applications may be required, 7-10 days apart.
Avoid spraying over flowering.

Fruit crops:

1.0 - 2.0L/ha. Post harvest, pre flowering
and fruit fill or as required.

Vegetables: 1.0 - 2.0L/ha. As required.

Vine crops:

1.0 - 2.0L/ha
Pre-flower, post fruit set and post harvest.

Broadacre crops:

1.0 - 2.0L/ha post emergence or as required.

Never spray leaves with more than 4% of solution. For example
for every 100lt of water you can apply 4lt of liquid fertilisers.



PROUDLY MADE IN AUSTRALIA



FOLIAR



20L



200L



1000L

